

Brent Island Meadow Survey. The abundance of forbs ('wild flowers'), grasses and other plants in the Lower (formerly Areas 1 & 2) and Upper (Area 3) in 2015, 2016 and 2017 using the Domin scale. The abundance given is the maximum recorded in either of the two surveys carried out each year - Early (early June) and Late (late July).

Overall occurrence			Species	Common name	2015		2016		2017	
2015	2016	2017			LOWER	UPPER	LOWER	UPPER	LOWER	UPPER
Forbs										
♦	♦	♦	<i>Achillea millefolium</i>	Yarrow	O	O	R	R	O	O
♦			<i>Aegopodium podagraria</i>	Ground-elder	R					
	♦	♦	<i>Angelica sylvestris</i>	Angelica			R		R	R
♦	♦	♦	<i>Centaurea nigra</i>	Common Knapweed	LF	O	LF	O	F	F
♦	♦	♦	<i>Cerastium fontanum</i>	Common Mouse-ear	R	O	O	R	O	O
♦	♦	♦	<i>Cirsium arvense</i>	Creeping Thistle		O		LF		LF
♦	♦	♦	<i>Conopodium majus</i>	Pignut	LF	F	LF	F	F	R
	♦	♦	<i>Dactylorhiza praetermissa</i>	Southern Marsh-orchid				R		R
♦	♦	♦	<i>Heracleum sphondylium</i>	Hogweed	LA	F	LA	F	F	F
	♦		<i>Hyacinthoides non-scriptus</i>	Bluebell			R			
♦	♦	♦	<i>Hypochaeris radicata</i>	Cat's-ear	LF	O	LF	O	F	LF
	♦	♦	<i>Leucanthemum vulgare</i>	Oxeye Daisy			R		R	
♦	♦	♦	<i>Lotus pedunculatus</i>	Greater Bird's-foot-trefoil	LF	O	LF	O	LF	
♦	♦	♦	<i>Oenanthe crocata</i>	Hemlock Water-dropwort	R	R		R		R
♦	♦	♦	<i>Plantago lanceolata</i>	Ribwort Plantain	F	A	F	F	A	A
♦	♦	♦	<i>Plantago major</i>	Greater Plantain	LF	LA	LF	LA	LA	LA
	♦	♦	<i>Ranunculus acris</i>	Meadow Buttercup			R	O		O
♦	♦	♦	<i>Ranunculus repens</i>	Creeping Buttercup	F	F	F	F	LA	A
	♦	♦	<i>Rhinanthus minor</i>	Yellow-rattle			LF	LF	LF	LF
♦	♦	♦	<i>Rumex acetosa</i>	Common Sorrel	F	F	F	F	A	F
♦	♦	♦	<i>Rumex obtusifolius</i>	Broad-leaved Dock	O			O	O	R
	♦	♦	<i>Rumex sanguineus</i>	Wood Dock			O	R	R	O
♦	♦	♦	<i>Stellaria graminea</i>	Lesser Stitchwort	O	O	O	R	LF	O
	♦	♦	<i>Taraxacum</i> agg.	Dandelion			R	R	R	R
♦	♦	♦	<i>Trifolium pratense</i>	Red Clover	O	F	O	F	O	F
♦	♦	♦	<i>Trifolium repens</i>	White Clover	LA	A	LA	A	A	A
♦	♦	♦	<i>Urtica dioica</i>	Common Nettle	R	O		LF		LF
♦	♦	♦	<i>Veronica chamaedrys</i>	Germander Speedwell	F	F	LF	F	F	F
Grasses										
♦	♦	♦	<i>Agrostis capillaris</i>	Common Bent	A	A	A	A	A	A
♦			<i>Alopecurus pratense</i>	Meadow Foxtail	R					
♦	♦	♦	<i>Anthoxanthum odoratum</i>	Sweet Vernal-grass	A	A	A	A	A	A
♦	♦	♦	<i>Arrhenatherum elatius</i>	False Oat-grass	O		O		LF	
♦	♦	♦	<i>Cynosurus cristatus</i>	Crested Dog's-tail	O	F	O	F	O	F
♦	♦	♦	<i>Dactylis glomerata</i>	Cock's-foot	F	F	F	O	F	F
♦	♦	♦	<i>Holcus lanatus</i>	Yorkshire-fog	A	A	A	A	A	F
♦	♦	♦	<i>Holcus mollis</i>	Creeping Soft-grass	F	A	LA	A	A	A
♦	♦	♦	<i>Lolium perenne</i>	Perennial Rye-grass	LF	LA	LF	LF	LA	LA
♦	♦	♦	<i>Phleum pratense</i>	Timothy	F	F	F	F	F	F
♦	♦	♦	<i>Poa annua</i>	Annual Meadow-grass	LF	LF	LF	LF	F	F
♦	♦	♦	<i>Poa trivialis</i>	Rough Meadow-grass	LF	LF	F	LF	F	A
Other relevant										
♦	♦	♦	<i>Luzula campestris</i>	Field Wood-rush	LF		LF		O	
♦	♦	♦	<i>Pteridium aquilinum</i>	Bracken	LA		LA	R	LF	
32	37		TOTAL NUMBER OF SPECIES		33	28	35	35	34	34